

AD-A281 613



①

OFFICE OF NAVAL RESEARCH

GRANT N00014-91-J-1784

R&T CODE 413v002

Technical Report No. 4

EXTRACELLULAR IONIC COMPOSITION ALTERS KINETICS OF VESICULAR
RELEASE OF CATECHOLAMINES AND QUANTAL SIZE DURING EXOCYTOSIS AT
ADRENAL MEDULLARY CELLS

by

Jeffrey A. Jankowski, Jennifer M. Finnegan, and R. Mark Wightman

Prepared for Publication in the
Journal of Neurochemistry

Department of Chemistry
University of North Carolina at Chapel Hill
Chapel Hill, NC 27599-3290

July 5, 1994

Reproduction in whole or in part is permitted
for any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC QUALITY INSPECTED 5

94 7 12 093

DTIC
ELECTE
JUL 14 1994
S-B-D

94-21509

33px

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302 and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 5, 1994		3. REPORT TYPE AND DATES COVERED Technical - 6/1993 - 6/1994	
4. TITLE AND SUBTITLE Extracellular Ionic Composition Alters Kinetics of Vesicular Release of Catecholamines and Quantal Size During Exocytosis at Adrenal Medullary Cells				5. FUNDING NUMBERS N00014-91-J-1784	
6. AUTHOR(S) J.A. Jankowski, J.M. Finnegan, and R.M. Wightman					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of North Carolina at Chapel Hill Chapel Hill, NC 27599-3290				8. PERFORMING ORGANIZATION REPORT NUMBER ONR Technical Report IV	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Submitted to Journal of Neurochemistry					
12a. DISTRIBUTION / AVAILABILITY STATEMENT This document has been approved for public release and sale; its distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The temporal resolution of carbon-fiber microelectrodes has been exploited to examine the plasticity of quantal secretory events at individual adrenal medullary cells. The size of individual quantal events, monitored by amperometric oxidation of released catecholamines, was found to be dependent on the extracellular ionic composition, the secretagogue, and the order of depolarization delivery. Release was observed with either exposure to 60 mM K ⁺ in the presence of Ca ²⁺ or exposure to 3 mM Ba ²⁺ in solutions of different pH, with and without external Ca ²⁺ . Ba ²⁺ was demonstrated to induce Ca ²⁺ -independent exocytotic release for an extended period of time (>4 min) relative to release induced by K ⁺ (~30 s), which is Ca ²⁺ -dependent. In all cases, simultaneous changes of intracellular divalent cations, monitored by fura-2 fluorescence, accompanied quantal release and had a similar time course. Exocytosis caused by Ba ²⁺ in Ca ²⁺ -free medium had a larger mean spike area at pH 8.2 than at pH 7.4. When Ba ²⁺ -induced spikes measured at pH 7.4 were compared, the spikes in Ca ²⁺ -free medium were found to be broader and shorter, but had the same area. Release induced by K ⁺ after exposure to Ba ²⁺ was comprised of larger quantal events when compared to preceding K ⁺ stimulations. Finally, spikes obtained with Ba ²⁺ exposure at an extracellular pH of 5.5 had a different shape than those obtained in more basic solutions. These changes in spike size and shape are consistent with the interactions between catecholamines and other intravesicular components.					
14. SUBJECT TERMS				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
20. LIMITATION OF ABSTRACT					

EXTRACELLULAR IONIC COMPOSITION ALTERS KINETICS OF VESICULAR
RELEASE OF CATECHOLAMINES AND QUANTAL SIZE DURING EXOCYTOSIS AT ADRENAL
MEDULLARY CELLS

Jeffrey A. Jankowski¹, Jennifer M. Finnegan, and R. Mark Wightman^{*}

Department of Chemistry
University of North Carolina at Chapel Hill
Chapel Hill, NC 27599-3290

^{*}To whom correspondence should be addressed.

¹Present address: University of Illinois at Urbana-Champaign
Neuronal Pattern Analysis Group
Beckman Institute
405 N. Matthews
Urbana, IL 61801

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

ABSTRACT

The temporal resolution of carbon-fiber microelectrodes has been exploited to examine the plasticity of quantal secretory events at individual adrenal medullary cells. The size of individual quantal events, monitored by amperometric oxidation of released catecholamines, was found to be dependent on the extracellular ionic composition, the secretagogue, and the order of depolarization delivery. Release was observed with either exposure to 60 mM K^+ in the presence of Ca^{2+} or exposure to 3 mM Ba^{2+} in solutions of different pH, with and without external Ca^{2+} . Ba^{2+} was demonstrated to induce Ca^{2+} -independent exocytotic release for an extended period of time (>4 min) relative to release induced by K^+ (~ 30 s), which is Ca^{2+} -dependent. In all cases, simultaneous changes of intracellular divalent cations, monitored by fura-2 fluorescence, accompanied quantal release and had a similar time course.

Exocytosis caused by Ba^{2+} in Ca^{2+} -free medium had a larger mean spike area at pH 8.2 than at pH 7.4. When Ba^{2+} -induced spikes measured at pH 7.4 were compared, the spikes in Ca^{2+} -free medium were found to be broader and shorter, but had the same area. Release induced by K^+ after exposure to Ba^{2+} was comprised of larger quantal events when compared to preceding K^+ stimulations. Finally, spikes obtained with Ba^{2+} exposure at an extracellular pH of 5.5 had a different shape than those obtained in more basic solutions. These changes in spike size and shape are consistent with the interactions between catecholamines and other intravesicular components.

Key words: chromaffin cells, catecholamine, barium induced exocytosis, fura-2, calcium independent exocytosis

Running title: Effects of extracellular ions on quantal release

INTRODUCTION

Many cells secrete chemical substances by exocytosis, whereby intracellular vesicles fuse with the plasma membrane and extrude their contents into the extracellular space (for reviews see Almers, 1990; DeCamilli and Jahn, 1990; Burgoyne and Morgan, 1993). This process has classically been viewed as an all-or-none secretory event (Martin, 1966; Viveros et al., 1969) and, as a result, is often referred to as quantal secretion (Bekkers et al., 1990). Single exocytotic events involving the release of electroactive substances can now be quantitated by oxidation with a carbon-fiber microelectrode placed adjacent to a cell (Leszczyszyn et al., 1990; Wightman et al., 1991; Chow et al., 1992; Schroeder et al., 1992; Alvarez de Toledo et al., 1993). This technique is used here to probe changes in exocytosis as indicated by the size and shape of individual current spikes.

Bovine adrenal medullary cells, which are used in this work, have played a benchmark role in numerous investigations of exocytosis (for reviews see Viveros, 1975; Pollard et al., 1985; Burgoyne, 1991). These cells secrete catecholamines which are stored in vesicles at a high concentration (>300 mM (Winkler, 1976; Schroeder, 1992)). The vesicles contain several other substances including Ca^{2+} (30 mM (Bulenda and Gratzl, 1985)) and chromogranin A (CGA, 1 mM). CGA is a highly acidic protein (Yoo and Albanesi, 1991; Helle et al., 1993; Winkler, 1976) which aggregates and binds Ca^{2+} and catecholamines at the intravesicular pH of 5.5 (Videen et al., 1992; Yoo and Lewis, 1992). As a result it has been proposed to play the role of a vesicle condensing protein (Simon and Aunis, 1989; Winkler and Fisher-Colbrie, 1992). Thus, exocytotic release must involve dissociation of the intravesicular contents as well as extrusion of the vesicle contents.

In previous work with bovine adrenal medullary cells, we have examined the characteristics of catecholamine concentration spikes resulting from individual exocytotic events induced by Ca^{2+} entry after digitonin permeabilization (Jankowski et al., 1993; Ciolkowski et al., submitted). Spike symmetry, amplitude, area, and frequency were found to be dependent on

the extracellular pH. Furthermore, at each pH value examined, the catecholamine spikes were temporally broadened relative to that expected for free diffusion of a concentration packet of catecholamines away from the cell surface (Schroeder et al., 1992). Partial release of vesicular contents has been detected with amperometric measurements of serotonin release from mast cells (Alvarez de Toledo et al., 1993). All of these observations suggest that the state of association of the vesicle contents can affect the temporal characteristics and quantal size of an exocytotic event.

In this work we investigate the shape of the amperometric spikes induced by exposure to Ba^{2+} . This secretagogue is unique in its ability to cause Ca^{2+} -independent secretion, and, therefore, allows examination of the effects of alteration of extracellular Ca^{2+} gradients on spike shape. The results show that the quantal size and time course of extrusion of the vesicular contents are affected both by the external pH and the external Ca^{2+} concentration.

EXPERIMENTAL PROCEDURES

Electrodes and Electrochemical Procedures. Carbon fibers (5- μ m radius, Thornell P-55, Amoco, Corp., Greenville, SC) were used to prepare microelectrodes (Kawagoe et al., 1993) which were polished to obtain an elliptical sensing surface with an average radii of 6 μ m. Electrodes were operated amperometrically with an applied potential of 0.650 V versus a saline-saturated calomel electrode.

Single Cell Stimulus-Secretion Experiments. Primary cultures of bovine adrenal medullary cells were prepared from fresh tissue and cultured as described previously (Wightman et al., 1991). Cell populations were enriched in epinephrine-containing cells by differential centrifugation with a single step Renografin gradient (Moro et al., 1990). Experiments were conducted within days 3 and 10 of culture. For release studies, the culture medium was replaced with the incubation buffer which contained 150 mM NaCl, 4.2 mM KCl, 1.0 mM Na_2HPO_4 , 11.2 mM glucose, 0.7 mM MgCl_2 , and 10 mM HEPES with the pH adjusted to the desired value using NaOH. In Ca^{2+} -containing solutions 3.0 mM CaCl_2 was employed. Without added Ca^{2+} , the buffer contained 2.8 μ M free Ca^{2+} as determined with an ion-selective electrode (Orion Model 93-20, Cambridge, MA). To further lower the concentration of free Ca^{2+} , 0.2 mM EGTA (with 6 mM NaCl to maintain osmolarity) was added to prepare the solutions termed Ca^{2+} -free. This was done at all pH values even though EGTA has a much lower affinity for Ca^{2+} at pH 5.5. BaCl_2 (3.0 mM in incubation buffer without Ca^{2+} or Mg^{2+}) and KCl (60 mM in incubation buffer with 3 mM CaCl_2 and NaCl reduced to 94.2 mM) were applied to cells via pressure ejection from micropipettes. Experiments were conducted at room temperature on the stage of an inverted microscope (Axiovert 35, Zeiss, Thornwood, NY). Carbon-fiber microelectrodes were positioned 1 μ m from the surface of individual cells using a piezoelectric driver (PCS-250 Patch Clamp Driver, Burleigh Instruments, Fishers, NY) as previously described (Schroeder et al, 1992).

Data Analysis. Unfiltered electrode responses were recorded on 1/2" videotape. When replayed, the records were low pass filtered at 2600 Hz

(CyberAmp 320, Axon Instruments, Inc., Foster City, CA), and computer digitized at 0.2 msec/pt (Fetchex, Axon Instruments, Foster City, CA). Spikes were computer selected with locally written software (Schroeder et al., 1992) if their amplitude was five times greater than the root-mean-square noise of the record. Skew values of the spikes were calculated as described by Foley and Dorsey (1983). Because the data is inherently nonparametric, significance was evaluated by the Mann-Whitney test (Systat, Evanston, IL).

Fluorescence studies. Chromaffin cells cultured on glass coverslips (Assistant Glass, Carolina Biological Supply, Burlington, NC) were loaded at room temperature for 60 minutes with 1 μ M fura-2 AM (Molecular Probes, Eugene, OR) in Krebs-Ringer buffer (145 mM NaCl, 5 mM KCl, 1.2 mM Na_2HPO_4 , 10 mM glucose, 1.3 mM MgCl_2 , and 20 mM HEPES, pH 7.4 (O'Sullivan et al., 1989)) containing 4 mM EGTA, and 0.1% bovine serum albumin. The culture plate was then rinsed twice and refilled with Krebs-Ringer buffer of the desired pH and Ca^{2+} concentration. Ionic concentrations were adjusted as in the stimulus-secretion studies. Fura-2 fluorescence measurements at single cells were made at room temperature with a 40x oil immersion objective (Fluor 40/1.3, Nikon, Melville, NY) on an inverted microscope equipped with an EMPIX Photometer System (EMPIX Imaging, Mississauga, Canada) which employs a 510 nm (40 nm bandpass) emission filter. Fluorescence intensities from alternate excitation at 340 nm and 380 nm were measured every 110 ms through a 27 μ m pinhole aperture at the photomultiplier tube, corrected for autofluorescence and electrode reflectance, and then ratioed (F_{340}/F_{380}).

Fura-2 can bind to Ba^{2+} as well as Ca^{2+} but with a greater affinity for Ca^{2+} than Ba^{2+} ($K_{d,\text{Ca}} = 224$ nM (Grynkiewicz et al., 1985), $K_{d,\text{Ba}} = 780$ nM (Schilling et al., 1989), both at 37°C). In addition, the wavelength for maximal excitation of fluorescence differ for the bound forms. This results in a difference in the maximum fluorescence ratio as measured both *in vitro* and *in situ* with our instrument. *In vitro* calibration curves were constructed in EGTA-buffered solutions which mimic the cytosol (115 mM KCl, 20 mM NaCl, 1 mM MgCl_2 , 10 mM KMOPS, pH 7.1 (Schilling et al., 1989)). The maximum ratio

(obtained with 5 mM free Ca^{2+}) was 8.68 times greater than that for Ba^{2+} (3 mM free concentration). *In situ* calibrations, obtained in single cells permeabilized with 2 μM ionomycin (Williams and Fay, 1990) showed some cell-to-cell variability, and the relative sensitivity to the two ions was only 65% of that found *in vitro*.

Reagents. Unless noted above, chemicals were obtained from Sigma (St. Louis, MO). Solutions were prepared in doubly distilled water.

RESULTS

Comparison of release caused by K^+ and Ba^{2+} at pH 7.4. At pH 7.4, a 3-s pressure ejection of 60 mM K^+ with 3 mM Ca^{2+} onto a single cell caused amperometric spikes (exocytotic release of catecholamine) with or without Ca^{2+} in the external media (Figure 1A,B). Simultaneous fluorescence measurements in cells preloaded with fura-2 showed a corresponding increase in the fluorescence ratio (increase in $[Ca^{2+}]_i$). Both responses returned to baseline values 30 to 60 s after the initial exposure. Exposure of individual cells to K^+ without Ca^{2+} did not cause amperometric spikes (Wightman et al., 1991) or a change in the fluorescence ratio.

Having established cell viability with two exposures to K^+ with Ca^{2+} , the same cells were exposed to 3.0 mM Ba^{2+} for 15 s. Amperometric spikes were observed whether or not Ca^{2+} was present in the external medium (Figure 1). Secretion lasted for at least 4 min and was accompanied by a sustained increase in the fluorescence ratio. The extended time course of both responses is in sharp contrast to the responses induced by 3-s exposure to 60 mM K^+ . (K^+ application for 15 s also led to transient release whereas 3-s applications of Ba^{2+} caused release for several minutes (data not shown)).

A short time (1-2 min) after the initial rise in the fluorescence ratio and release of catecholamine following exposure of the cells to Ba^{2+} , concomitant fluctuations in the ratio and release were consistently observed at cells in Ca^{2+} -containing buffer (Figure 1B). Such cytosolic Ca^{2+} oscillations are often observed during long-lasting stimuli which alter internal Ca^{2+} stores (Berridge, 1990; Tse et al., 1993).

For experiments in media containing Ca^{2+} the cells were exposed to K^+ (3-s pressure ejection, two exposures) after the Ba^{2+} -induced release had subsided. The fluorescence ratio increased with K^+ as observed before exposure to Ba^{2+} . However, the characteristics of the spikes were altered relative to those induced by K^+ at the same cell before exposure to Ba^{2+} (Table 1). The means for the spike area (Q) and width at half-height ($\tau_{1/2}$) were significantly increased.

Spike characteristics as a function of pH. Amperometric spikes were also induced by Ba^{2+} from cells in buffers at pH 8.2 (Figure 2) or 5.5 (Figure 1C,D) whether or not Ca^{2+} was present in the buffer. In contrast, K^+ at pH 5.5 is unable to induce release although small changes in the fluorescent ratio were seen when the buffer contained Ca^{2+} (Figure 1D). The average characteristics (area, width at half height, and peak current (i_{max})) for Ba^{2+} induced spikes are summarized in Table 2. The spike area is particularly significant because it is a direct measure of the quantity of material detected, and thus, the mean value is the quantal size (Jankowski et al., 1992). Under all conditions, a broad distribution of Q values is observed (e.g., Figure 2B). This distribution is consistent with the variation in chromaffin cell vesicular radii (Schroeder et al., 1992).

The experimental results with the most significant differences are summarized in the following sections.

Ba^{2+} -induced spikes at pH 7.4 and 8.2 in the absence of Ca^{2+} . Release at pH 7.4 and 8.2 appears qualitatively similar (Figure 2A). However, the distribution of spike areas (Figure 2B) and their mean values (Table 2) show that the quantal size is increased at pH 8.2. In Ca^{2+} -free medium, the mean quantal size of spikes induced by Ba^{2+} is 50% greater at pH 8.2 than at pH 7.4, a significant ($p < 0.001$) change seen as a shift in the histogram.

The two spikes in Figure 2C have the average values obtained at each pH. At pH 8.2 the mean $t_{1/2}$ value is increased by 30% relative to that at pH 7.4, a significant ($p < 0.001$) change. The increase in $t_{1/2}$, which reflects the rates of dissociation and diffusion of the vesicular contents to the microelectrode surface, demonstrates that the increase in quantal amount is accompanied by an alteration in the rate of vesicular extrusion.

Ba^{2+} -induced spikes at pH 7.4 as a function of external Ca^{2+} . The mean quantal sizes following exposure to Ba^{2+} at pH 7.4 are the same regardless of the extracellular Ca^{2+} concentration (Table 2). However, in the absence of Ca^{2+} the mean amplitude of spikes induced by Ba^{2+} are significantly ($p < 0.001$) lower and the mean half widths are significantly ($p < 0.001$) wider than those

obtained in medium containing Ca^{2+} (37% decrease in mean i_{max} , 50% increase in mean $t_{1/2}$). Histograms of these values clearly show these differences (Figure 3A). When Ca^{2+} is absent from the buffer, fewer large amplitude spikes and fewer narrow spikes are observed. Spikes which have the mean characteristics of those measured during exposure to Ba^{2+} at pH 7.4 are shown in Figure 3B for solutions with and without Ca^{2+} .

Spike shape at pH 5.5. Under physiological conditions, the shape of individual spikes has been found to be similar to that predicted for diffusional dispersion from a point source (Schroeder et al., 1992; Jankowski et al., 1993; Ciolkowski et al., submitted). The shape of such a spike is characterized by a sharp rise followed by slow decay, resembling an exponentially modified gaussian (Schroeder et al., 1992). Qualitatively, a significant number of the spikes obtained in response to Ba^{2+} at pH 5.5 without extracellular Ca^{2+} appeared to be more symmetric than expected for diffusional dispersion.

To quantitatively analyze their symmetry, skew values for the spikes were determined. The measure of skew employed was originally developed to characterize the asymmetry of chromatographic peaks (Foley and Dorsey, 1983). In this analysis, a diffusion-controlled shape has a skew value (termed v_s) of 1.8 while a purely gaussian shape has a value of 0.4. Because the method is based on the peak width at 10% of the peak height, skew values less than 1.7 also may indicate the presence of a pre-spike feature (Chow et al., 1992). Example spikes are shown in Figure 4. The spike in Figure 4A has a skew value of 0.41, and noticeably deviates from the superimposed diffusion-controlled shape. The spike in Figure 4B has a low skew value (0.77) caused by a pre-spike feature. Figure 4C shows a spike with a skew of 1.0, and in Figure 4D a spike is shown with a skew of 1.8 which correlates ($r = 0.98$) with that predicted for a diffusion-controlled shape. As these examples show, the spikes with low skew values have a greater shape symmetry or a distinct pre-spike feature.

Skew values of spikes detected at pH 5.5 were compared to those obtained

at pH 8.2 (Jankowski et al., 1993). At pH 8.2 in the absence of Ca^{2+} , the mean skew value is 1.78 and 96.7% of the spikes fall in the range of 1.7-1.9. However, at pH 5.5 without Ca^{2+} , 30.5% of the spikes have skew values less than 1.7 indicating a deviation from the diffusion-controlled shape.

DISCUSSION

In this study we establish that the release of catecholamines induced by Ba^{2+} , both in the absence and presence of Ca^{2+} , occurs as a prolonged series of sharp, amperometric spikes. This confirms that an influx of Ba^{2+} can substitute for Ca^{2+} as a trigger for exocytosis. Second, we show that the characteristics of the packets of catecholamine released by Ba^{2+} are dependent on the composition of the extracellular medium. A hypothesis which accounts for this observation is that the temporal characteristics of the individual vesicular events are dependent on the state of association of the vesicle contents prior to exocytosis.

In prior work we have shown that the area and distribution of amperometric spikes at pH 7.4 is independent of the secretagogue employed and its concentration (Wightman et al., 1991). Release induced by Ba^{2+} in the presence of Ca^{2+} at pH 7.4 shares this characteristic in that the mean characteristics of the spikes are similar to those induced by K^+ (Tables 1 and 2). However, release induced by Ba^{2+} continues for a longer time than that induced by K^+ (Figure 1) even though both secretagogues are applied transiently in the form of small droplets. The droplets are rapidly diluted by the incubation medium as a result of diffusion and convection (Leszczyszyn et al., 1991) on a time scale that is approximately that for the release of catecholamine following exposure to K^+ . The Ba^{2+} -induced release continues for 3-15 minutes after its initial application even in medium containing EGTA which should dissipate it even more rapid because of complexation. This suggests Ba^{2+} is retained in the cell where it can promote release either directly, by mobilization of internal Ca^{2+} stores, or by blockade of Ca^{2+} efflux (Przywara et al., 1993).

Prior work has shown that Ba^{2+} is capable of inducing massive secretion from several cell types, including adrenal cells, that is independent of extracellular Ca^{2+} (Douglas and Rubin, 1964a; Heldman et al., 1989; Morita et al., 1990; TerBush and Holz, 1992; Przywara et al., 1993). Ba^{2+} may enter through both voltage- and receptor-gated Ca^{2+} channels (Forsberg and Pollard,

1988; Heldman and Pollard, 1989), and, once inside the cell, directly cause release of catecholamines (TerBush and Holz, 1992). Irrespective of the mechanism, the long-lasting and high-frequency occurrence of spikes following exposure to Ba^{2+} (Figure 1) establishes that Ba^{2+} can directly induce exocytosis, and can do so in the absence of extracellular Ca^{2+} .

Entry of Ca^{2+} into adrenal medullary cells is the normal trigger for exocytosis (Burgoyne, 1991). The simultaneous observations of intracellular Ca^{2+} elevation and catecholamine release in response to K^+ (Figure 1) supports a close temporal link between these two events at pH 7.4. At acidic pH values Na^+ channel activity is reduced (Rutten et al., 1989), and lower $[Ca^{2+}]_i$ changes are observed that are not accompanied by catecholamine release.

The fura-2 fluorescence measurements also show that extracellular Ba^{2+} causes an increase in intracellular divalent ions. While a portion of the measured signal is undoubtedly due to Ba^{2+} which enters the cell (Heldman et al., 1989, TerBush and Holz, 1992), the fluorescent ratio obtained in Ca^{2+} free media exceeds the $R_{max,Ba}$ obtained by *in situ* calibrations by 18 % ($n = 5$). Therefore, a portion of the increase must be due to liberated intracellular Ca^{2+} . However, since Ba^{2+} does not elevate the intracellular messenger IP_3 (TerBush and Holz, 1992) and has been suggested to have poor affinity for the Ca^{2+} -ATPase (Schilling, 1989), the intracellular source of Ca^{2+} does not appear to be the endoplasmic reticulum. A similar interpretation has been invoked to explain Ba^{2+} induced changes in cultured sympathetic neurons (Przywara et al., 1993). While this paper was being reviewed, fura-2 measurements were published which also indicate a contribution of both Ba^{2+} and Ca^{2+} to the Ba^{2+} -induced fluorescence ratios (von Rüden et al., 1993).

The most surprising results of these experiments is that the characteristics of spikes induced by Ba^{2+} can be altered by the external media. First, exocytosis caused by Ba^{2+} in Ca^{2+} -free medium has a larger mean spike area at pH 8.2 than at pH 7.4. The smaller mean "quanta" at pH 7.4 suggests that a significant percentage of catecholamine is either not released

upon vesicle fusion (Alvarez de Toledo et al., 1993), or is released at a very slow, undetectable rate. Second, when Ba^{2+} -induced spikes measured at pH 7.4 are compared, the areas of the spikes are independent of the presence of Ca^{2+} in the external medium, but the spikes in Ca^{2+} -free medium are broader and shorter. Furthermore, release induced by Ba^{2+} at pH 7.4 appears to alter the quantal size of spikes induced by subsequent exposure to K^+ . Third, while large area spikes are obtained at an extracellular pH of 5.5, the shape of some of the spikes is quite different from those in more basic solutions. Although some of the external media employed are extreme conditions for the cells, the consistently low basal fura-2 ratios at all pH values show that the cells retain integrity with respect to Ca^{2+} entry.

A possible reason for these differences in size and shape of the spikes is the extent of aggregation or association of the vesicular matrix before and even during exocytosis. This association depends in large part on the conformation of chromogranin-A, CGA, the primary water soluble protein of the vesicles in adrenal medullary cells (Winkler and Fisher-Colbrrie, 1992). Because CGA can bind Ca^{2+} (Reiffen and Gratzl, 1986; Gorr et al., 1988; Westermann et al., 1988; Yoo and Albanesi, 1990a & b; Yoo and Albanesi, 1991; Videen et al., 1992; Yoo and Lewis, 1992), the vesicles are a major storage site of Ca^{2+} in the adrenal medullary cell (Yoo and Albanesi, 1990b). Prior studies have shown that pH and Ca^{2+} affect the conformation of CGA and its ability to aggregate catecholamine (Gorr et al., 1988, Westermann et al., 1988; Yoo and Albanesi, 1990a; Yoo and Albanesi, 1991). At the intravesicular pH of 5.5, CGA exists predominantly as a tetramer with a high α -helical content, while at the physiological pH value normally encountered upon fusion, CGA is less aggregated and exists predominantly as a dimer with less α -helical character (Yoo and Lewis, 1992). CGA also has a lower binding capacity for Ca^{2+} and catecholamines at pH 7.4, although calculations suggest that even at pH 5.5 only 5.3% of the intravesicular catecholamine is chemically bound by CGA (Videen et al., 1992).

Thus, we hypothesize that the observed changes in spike characteristics

are due to the presence of concentration gradients of the intravesicular ions and molecules that are initially generated upon vesicle fusion which, when increased, promote further dissociation of the vesicular matrix. The condition which maximizes the quantity of material released by exocytosis, high pH and low Ca^{2+} , is that which maximizes the disaggregation of CGA (Gorr et al., 1988). Since the measured quantal size (Q) at pH 7.4 with or without Ca^{2+} never reaches the value found at pH 8.2, exocytosis induced by secretagogues at pH 7.4 must result in a partial or slowed release of vesicular contents. Continued attachment of some CGA to the vesicle int. (Yoo, 1993) would provide a matrix for withholding catecholamines. Alternatively, partial exocytosis may result from closure of the fusion pore (Alvarez de Toledo et al., 1993). The altered response to K^+ following exposure to Ba^{2+} at pH 7.4 may indicate an alteration of intravesicular aggregation caused by Ba^{2+} entry. However, we can not eliminate the possibility that, since Ba^{2+} causes prolonged release, the population of vesicles from which release occurs during the latter exposure to K^+ may be different from the original population.

The spikes at pH 8.2 in the absence of Ca^{2+} are wider than at other pH values, and much wider than expected for free diffusion of catecholamines (Jankowski et al., 1993). CGA could behave like other vesicular proteins present in different cells (Verdugo, 1990; Chandler, M. et al., 1989; Merkle and Chandler, 1991) and swell upon fusion. Swelling, with its associated liberation of catecholamines could be kinetically limiting. Alternatively, swollen CGA could cause slow diffusion of the catecholamines from the cell surface to the microelectrode because of interaction with the prevalent anionic sites in the swollen CGA matrix (Winkler and Smith, 1975). The effect of Ca^{2+} on spike width at pH 7.4 is also consistent with these interpretations. The presence of EGTA in the external medium may promote matrix swelling by increasing the Ca^{2+} concentration gradient. An alternate factor that could affect the spike width is that the fusion pore formed upon exocytosis restricts the rates and amounts of vesicular extrusion. However,

there is no evidence that formation of the fusion pore is pH-dependent. Indeed, the fusion pore in mast cells has been shown to form with equal probability at low and physiological pH values (Monck et al., 1991).

According to the above hypothesis, exocytosis into an extracellular medium of pH 5.5, which eliminates the pH gradient, would result in little disaggregation of the matrix. Indeed, we have shown previously that the frequency of exocytotic events are greatly diminished at this pH unless a large catecholamine concentration gradient is generated by the carbon-fiber microelectrode (Jankowski et al., 1993). The spikes at pH 5.5 also tend to be narrower than at more basic pH values, and have lower skew values than predicted for diffusional processes. These differences suggest another process becomes rate limiting such as a change from slow diffusion to slow matrix dissociation or perhaps ionic effects on cell membrane dynamics. While narrow, the spikes are still wider than predicted for diffusion of catecholamines in free solution (Jankowski et al., 1993).

In summary, these investigations suggest two different ways in which the kinetics of exocytosis can be affected. First, the quantity of the released packet of catecholamine can be affected by the concentration gradient of H^+ ions which occurs upon vesicular fusion. Alteration of the extracellular medium to favor disaggregation of the matrix increases the mean quantal size of catecholamine. Second, an increase of the Ca^{2+} and pH gradients which occur upon vesicle fusion, by adjustment of the external medium, can cause an increase in the time of dissociation of the vesicular matrix as seen in the increase in half width of the individual spikes.

ACKNOWLEDGEMENTS

This research was supported by the Office of Naval Research and a BRSG from NIH to the University of North Carolina at Chapel Hill. JMF was the recipient of a fellowship from the Department of Education.

FIGURE LEGENDS

Figure 1. Simultaneous measurement of fura-2 fluorescence ratio (upper traces) and amperometric release (lower traces) at single adrenal medullary cells. Arrows indicate pressure ejection of secretagogue (K^+ : 3 s application of 60 mM K^+ and 3 mM Ca^{2+} ; Ba^{2+} : 15 s application of 3 mM Ba^{2+}). (A) Cell in Krebs-Ringer buffer containing 0.2 mM EGTA, pH 7.4. (B) Cell in Krebs-Ringer buffer containing 3 mM Ca^{2+} , pH 7.4. (C) Cell in Krebs-Ringer buffer containing 0.2 mM EGTA, pH 5.5. (D) Cell in Krebs-Ringer buffer containing 3 mM Ca^{2+} , pH 5.5. Note that for *in vitro* calibrations, $R_{max,Ba} = 0.38$.

Figure 2. Catecholamine release from adrenal medullary cells induced by Ba^{2+} in Ca^{2+} -free media. (A) Amperometric traces recorded at two different cells at the pH values noted on the traces. At the time indicated by the bar, 3 mM Ba^{2+} was applied for 15 s by pressure ejection from a micropipette. (B) Histogram of the charge measured for individual spikes at the pH values indicated. Results are from 657 spikes collected at 5 cells at pH 7.4 and 413 spikes collected from 5 cells at pH 8.2. (C) A single spike from each of the pH values which has the mean characteristics of the data set (see Table 2). The definitions for i_{max} and $t_{1/2}$ are indicated on the spike recorded at pH 7.4.

Figure 3. Secretion of catecholamines induced by Ba^{2+} at pH 7.4 with and without extracellular Ca^{2+} . (A) Width at half height and peak current histograms. Filled bars are results from 757 spikes obtained at 5 cells with 3 mM Ca^{2+} in the medium. Open bars are results from 657 spikes obtained at 5 cells with 0.2 mM EGTA. (B) Representative spikes with the mean spike characteristics for the data sets (see Table 2) with (dashed line) and without (solid line) extracellular Ca^{2+} are superimposed.

Figure 4. Spikes detected in response to 3.0 mM Ba^{2+} at pH 5.5 with no extracellular Ca^{2+} . Superimposed on each spike (dashed line) is the exponentially modified gaussian shape for diffusional dispersion. (A) Skew = 0.41, $r = 0.85$. (B) Skew = 0.77, $r = 0.92$. (C) Skew = 1.0, $r = 0.84$. (D) Skew = 1.8, $r = 0.98$ (r is the correlation coefficient for the experimental data with the diffusion-based shape (Schroeder et al., 1992)).

TABLE 1. Mean characteristics of spikes measured at cells in pH 7.4 buffer with 3.0 mM extracellular Ca^{2+} stimulated for 3 s with 60 mM K^+ before and after a 15-s exposure to 3.0 mM Ba^{2+} . Spikes were measured with a 6- μm radius electrode placed 1 μm away from the cell surface. Data was collected at 200 $\mu\text{sec/pt}$ and low pass filtered at 2600 Hz. Means are reported \pm s.e.m.

Stimulus	60 mM K^+	
	Before Ba^{2+}	After Ba^{2+}
Q (pC)	1.17 \pm .07	1.58* \pm .09
$t_{1/2}$ (ms)	7.23 \pm .38	10.1* \pm .50
I_{max} (pA)	129 \pm 6.2	120 \pm 5.8
f (Hz)	1.22	1.72
# of spikes	305	276
# of cells	5	5

* Means are significantly different from 60 mM K^+ before Ba^{2+} exposure (Mann-Whitney test $p < 0.001$).

TABLE 2. Mean characteristics of spikes measured at cells exposed to 3.0 mM Ba^{2+} for 15 s in the presence and absence of 3.0 mM Ca^{2+} and at different pH values. Spikes were measured with a 6- μm radius electrode placed 1 μm from the cell surface. Data was collected at 200 $\mu\text{sec/pt}$ and low pass filtered at 2600 Hz. Means are reported \pm s.e.m.

pH	5.5		7.4		8.2	
$[\text{Ca}^{2+}]$	0 mM	3 mM	0 mM	3 mM	0 mM	3 mM
Q (pC)	1.87 ± 0.07	1.67 ± 0.07	1.35 ± 0.05	1.26 ± 0.05	2.02 ± 0.09	1.37 ± 0.10
$\tau_{1/2}$ (ms)	11.4 ± 0.38	11.0 ± 0.23	12.5 ± 0.30	8.34 ± 0.26	16.4 ± 0.54	12.7 ± 0.84
i_{max} (pA)	124 ± 4.5	95.2 ± 3.1	73.7 ± 2.2	117 ± 3.8	89.9 ± 3.7	85.7 ± 6.1
f (Hz)	0.78	0.97	1.14	1.22	0.61	0.77
# of spikes	557	419	657	757	413	173
# of cells	5	5	5	5	6	4

REFERENCES

- Almers, W. (1990) Exocytosis, *Annu. Rev. Physiol.* 52, 607-624.
- Alvarez de Toledo, G., Fernández-Chacón, R., and Fernández, J.-M. (1993) Release of Secretory Products During Transient Vesicle Fusion, *Nature* 363, 554-558.
- Bekkers, J. M., Richerson, G. B., and Stevens, C. F. (1990) Variability in Quantal Size in Cultured Hippocampal Neurons and Hippocampal Slices, *P. Natl. Acad. Sci.* 87, 5369-5362.
- Berridge, M.J. (1990) Calcium Oscillations, *J. Biol. Chem.* 265, 9583-9586.
- Bulenda, D. and Gratzl, M. (1985) Matrix free Ca^{2+} in Isolated Chromaffin Vesicles, *Biochemistry*, 24, 7760-7765.
- Burgoyne, R. D. (1991) Control of Exocytosis in Adrenal Chromaffin Cells, *Biochim. Biophys. Acta* 1071, 174-202.
- Burgoyne, R. D. and Morgan, A. (1993) Regulated Exocytosis, *Biochem J.* 293, 305-316.
- Chandler, D. E., Whitaker, M., and Zimmerberg, J. (1989) High Molecular Weight Polymers Block Cortical Granule Exocytosis in Sea Urchin Eggs at the Level of Granule Matrix Disassembly, *J. Cell Biol.* 109, 1269-1278.
- Chow, R. H., von Rüden, L., and Neher, E. (1992) Delay in Vesicle Fusion Revealed by Electrochemical Monitoring of Single Secretory Events in Adrenal Chromaffin Cells, *Nature* 356, 60-63.
- DeCamilli, P. and Jahn, R. (1990) Pathways to Regulated Exocytosis in Neurons, *Annu. Rev. Physiol.* 52, 625-645.
- Douglas, W. W. and Rubin, R. P. (1964a) Stimulant Action of Barium on the Adrenal Medulla, *Nature* 203, 305-307.
- Foley, J. P. and Dorsey, J. G. (1983) Equations for Calculation of Chromatographic Figures of Merit for Ideal and Skewed Peaks, *Anal. Chem.* 55, 730-737.
- Forsberg, E. J. and Pollard, H. B. (1988) Ba^{2+} -Induced ATP Release from Adrenal Medullary Chromaffin Cells is Mediated by Ba^{2+} -Entry Through Both Voltage- and Receptor-Gated Ca^{2+} Channels, *Neurosci.* 27, 711-715.

REFERENCES

- Almers, W. (1990) Exocytosis, *Annu. Rev. Physiol.* 52, 607-624.
- Alvarez de Toledo, G., Fernández-Chacón, R., and Fernández, J.-M. (1993) Release of Secretory Products During Transient Vesicle Fusion, *Nature* 363, 554-558.
- Bekkers, J. M., Richerson, G. B., and Stevens, C. F. (1990) Variability in Quantal Size in Cultured Hippocampal Neurons and Hippocampal Slices, *P. Natl. Acad. Sci.* 87, 5369-5362.
- Berridge, M.J. (1990) Calcium Oscillations, *J. Biol. Chem.* 265, 9583-9586.
- Bulenda, D. and Gratzl, M. (1985) Matrix free Ca^{2+} in Isolated Chromaffin Vesicles, *Biochemistry*, 24, 7760-7765.
- Burgoyne, R. D. (1991) Control of Exocytosis in Adrenal Chromaffin Cells, *Biochim. Biophys. Acta* 1071, 174-202.
- Burgoyne, R. D. and Morgan, A. (1993) Regulated Exocytosis, *Biochem J.* 293, 305-316.
- Chandler, D. E., Whitaker, M., and Zimmerberg, J. (1989) High Molecular Weight Polymers Block Cortical Granule Exocytosis in Sea Urchin Eggs at the Level of Granule Matrix Disassembly, *J. Cell Biol.* 109, 1269-1278.
- Chow, R. H., von Rüden, L., and Neher, E. (1992) Delay in Vesicle Fusion Revealed by Electrochemical Monitoring of Single Secretory Events in Adrenal Chromaffin Cells, *Nature* 356, 60-63.
- DeCamilli, P. and Jahn, R. (1990) Pathways to Regulated Exocytosis in Neurons, *Annu. Rev. Physiol.* 52, 625-645.
- Douglas, W. W. and Rubin, R. P. (1964a) Stimulant Action of Barium on the Adrenal Medulla, *Nature* 203, 305-307.
- Foley, J. P. and Dorsey, J. G. (1983) Equations for Calculation of Chromatographic Figures of Merit for Ideal and Skewed Peaks, *Anal. Chem.* 55, 730-737.
- Forsberg, E. J. and Pollard, H. B. (1988) Ba^{2+} -Induced ATP Release from Adrenal Medullary Chromaffin Cells is Mediated by Ba^{2+} -Entry Through Both Voltage- and Receptor-Gated Ca^{2+} Channels, *Neurosci.* 27, 711-715.

- Gorr, S.-U., Dean, W. L., Radley, T. L., and Cohn, D. V. (1988) Calcium-Binding and Aggregation Properties of Parathyroid Secretory Protein-I (Chromogranin A), *Bone and Mineral* 4, 17-25.
- Grynkewicz, G., Poenie, M., and Tsien, R. Y. (1985) A New Generation of Ca^{2+} Indicators with Greatly Improved Fluorescence Properties, *J. Biol. Chem.* 260, 3440-3450.
- Heldman, E., Levine, M., Raveh, L., and Pollard, H. B. (1989) Barium Ions Enter Chromaffin Cells via Voltage-Dependent Calcium Channels and Induce Secretion by a Mechanism Independent of Calcium, *J. Biol. Chem.* 264, 7914-7920.
- Helle, K. B., Serck-Hanssen, G., and Aardal, S. (1993) Functional Aspects of the Adrenal Medullary Chromogranins, *Neurochem. Int.* 22, 353-360.
- Jankowski, J. A., Schroeder, T. J., Holz, R. W., and Wightman, R. M. (1992) Quantal Secretion of Catecholamines Measured from Individual Bovine Adrenal Medullary Cells Permeabilized with Digitonin, *J. Biol. Chem.* 267, 18329-18335.
- Jankowski, J. A., Schroeder, T. J., Ciolkowski, E. L., and Wightman, R. M. (1993) Temporal Characteristics of Quantal Secretion of Catecholamines from Adrenal Medullary Cells, *J. Biol. Chem.* 268, 14694-14700.
- Kawagoe, K. T., Jankowski, J. A., and Wightman, R. M. (1991) Etched Carbon-Fiber Electrodes as Amperometric Detectors of Catecholamine Secretion from Isolated Biological Cells, *Anal. Chem.* 63, 1589-1593.
- Kawagoe, K. T., Zimmerman, J. B., and Wightman, R. M. (1993) Principles of voltammetry and microelectrode surface states, *J. Neurosci. Methods* 48, 225-240.
- Leszczyszyn, D. J., Jankowski, J. A., Viveros, O. H., Diliberto, E. J., Jr., Near, J. A., and Wightman, R. M. (1990) Nicotinic-Receptor Mediated Catecholamine Secretion from Individual Chromaffin Cells: Chemical Evidence for Exocytosis, *J. Biol. Chem.* 265, 14736-14737.
- Martin, A. R. (1966) Quantal Nature of Synaptic Transmission, *Physiol. Rev.* 46, 51-66.

- Merkle, C. J. and Chandler, D. E. (1991) Cortical Granule Matrix Disassembly during Exocytosis in Sea Urchin Eggs, *Develop. Biol.* 148, 429-441.
- Monck, J. R., Alvarez de Toledo, G., and Fernández, J. M. (1990) Tension in the Secretory Granule Membranes Causes Extensive Membrane Transfer Through the Exocytosis, *Proc. Natl. Acad. Sci., USA* 87, 7804-7808.
- Monck, J. R., Oberhauser, A. F., Alvarez de Toledo, G., and Fernández, J.-M. (1991) Is Swelling of the Secretory Granule Matrix the Force That Dilates the Exocytotic Fusion Pore?, *Biophys. J.* 59, 39-47.
- Morita, K., Teraoka, K., Azuma, M., Oka, M., and Hamano, S. (1990) Stimulatory Action of Ba^{2+} on Catecholamine Biosynthesis in Cultured Bovine Adrenal Chromaffin Cells: Possible Relation to Protein Kinase C, *Japan. J. Pharmacol.* 54, 425-432.
- Moro, M. A., Lopez, M. G., Gandia, L., Michelena, P., and Garcia, A. G. (1990) Separation and Culture of Living Adrenaline- and Noradrenaline-Containing Cells from Bovine Adrenal Medullae, *Anal. Biochem.* 185, 243-248.
- O'Sullivan, A. J., Cheek, T. R., Moreton, R. B., Berridge, M. J., and Burgoyne, R. D. (1989) Localization and heterogeneity of agonist-induced changes in cytosolic calcium concentration in single bovine adrenal chromaffin cells from video imaging of fura-2, *EMBO J.* 8, 401-411.
- Pollard, H. B., Ornberg, R., Levine, M., Kelner, K., Morita, K., Levine, R., Forsberg, E., Brocklehurst, K. W., Duong, L., Lelkes, P. I., Heldman, E., and Youdim, M. (1985) Hormone Secretion by Exocytosis with Emphasis on Information from the Chromaffin Cell System, *Vitam. Horm.* 42, 109-196.
- Przywara, D. A., Chowdhury, P. S., Bhave, S. V., Wakade, T. D., and Wakade, A. R. (1993) Barium-Induced Exocytosis is Due to Internal Calcium Release and Block of Calcium Efflux, *Proc. Natl. Acad. Sci., USA* 90, 557-561.
- Reiffen, F. U. and Gratzl, M. (1986) Ca^{2+} Binding to Chromaffin Vesicle Matrix Proteins: Effect of pH, Mg^{2+} , and Ionic Strength, *Biochem.* 25, 4402-4406.
- Ruffen, M.J., Delcore, R., Soybel, D.I., Moore, C.D. and Cheung, L.Y. (1989) Effects of cations and pH on apical membrane potential of in vitro *Necturus antrum*, *Am. J. Physiol.* 256, G798-807.

- Schilling, W. P., Rajan, L., and Strobl-Jager, E. (1989) Characterization of the Bradykinin-Stimulated Calcium Influx Pathway of Cultured Vascular Endothelial Cells, *J. Biol. Chem.* 264, 12838-12848.
- Schroeder, T. J., Jankowski, J. A., Kawagoe, K. T., Wightman, R. M., Lefrou, C., and Amatore, C. (1992) Analysis of Diffusional Broadening of Vesicular Packets of Catecholamine Released from Biological Cells During Exocytosis, *Anal. Chem.* 64, 3077-3083.
- Simon, J.-P. and Aunis, D. (1989) Biochemistry of the Chromogranin A Protein Family, *Biochem. J.* 262, 1-13.
- TerBush, D. R. and Holz, R. W. (1992) Barium and Calcium Stimulate Secretion from Digitonin-Permeabilized Bovine Adrenal Chromaffin Cells by Similar Pathways, *J. Neurochem.* 58, 680-687.
- Tse, A., Tse, F. W., Almers, W., and Hille, B. (1993) Rythmic Exocytosis Stimulated by GnRH-Induced Calcium Oscillations in Rat Gonadotropes, *Science* 260, 82-84.
- Verdugo, P. (1990) Goblet Cells Secretion and Mucogenesis, *Annu. Rev. Physiol.* 52, 157-176.
- Videen, J. S., Mezger, M. S., Chang, Y.-M., and O'Connor, D. T. (1992) Calcium and Catecholamine Interactions with Adrenal Chromogranins: Comparison of Driving Forces in Binding and Aggregation, *J. Biol. Chem.* 267, 3066-3073.
- Viveros, O. H., Arqueros, L., and Kirshner, N. (1969) Quantal Secretion from the Adrenal Medulla All-or-None Release of Storage Vesicle Content, *Science* 165, 911-913.
- Viveros, O. H. (1975) Mechanisms of secretion of catecholamines from adrenal medulla, in *Handbook of Physiology, Endocrinology*, Vol. 6, (H. Blaschko, G. Sayers, and A. D. Smith, eds.), pp. 389-426, American Physiological Society, Washington, DC.
- von Rüden, L., Garcia, A.G., and Lopez, M.G. (1993) The mechanism of Ba^{2+} -induced exocytosis from single chromaffin cells, *FEBS Lett.* 336, 48-52.

- Westermann, R., Stögbauer, K., Unsicker, K., and Lietzke, R. (1988) Calcium-Dependence of Chromogranin A-Catecholamine Interaction, *FEBS Lett.* 239, 203-206.
- Wightman, R. M., Jankowski, J. A., Kennedy, R. T., Kawagoe, K. T., Schroeder, T. J., Leszczyszyn, D. J., Near, J. A., Diliberto, E. J., Jr., and Viveros, O. H. (1991) Temporally Resolved Catecholamine Spikes Correspond to Single Vesicle Release From Individual Chromaffin Cells, *Proc. Natl. Acad. Sci.* 88, 10754-10758.
- Williams, D.A. and Fay, F.S. (1990) Intracellular calibration of the fluorescent calcium indicator Fura-2, *Cell Calcium* 11, 75-83.
- Winkler, H. (1976) The Composition of Adrenal Chromaffin Granules: An Assessment of Controversial Results, *Neurosci.* 1, 65-80.
- Winkler, H. and Smith, A. D. (1975) The Chromaffin Granule and the Storage of Catecholamines, in *Handbook of Physiology, Endocrinology*, Vol. 6, (H. Blaschko, G. Sayers, and A. D. Smith, eds.), pp. 367-388, American Physiological Society, Washington, DC.
- Winkler, H. and Fisher-Colbrrie, R. (1992) The Chromogranins A and B: The First 25 Years and Future Perspectives, *Neurosci.* 5, 1803-1823.
- Yoo, S. H. (1993) pH-Dependent Association of Chromogranin A with Secretory Vesicle Membrane and a Putative Membrane Binding Region of Chromogranin A, *Biochem.* 32, 8213-8219.
- Yoo, S. H. and Albanesi, J. P. (1990a) Inositol 1,4,5-trisphosphate-Triggered Ca^{2+} Release from Bovine Adrenal Medullary Secretory Vesicles, *J. Biol. Chem.* 265, 13446-13448.
- Yoo, S. H. and Albanesi, J. P. (1990b) Ca^{2+} -Induced Conformational Change Aggregation of Chromogranin A, *J. Biol. Chem.* 265, 14414-14421.

- Yoo, S. H. and Lewis, M. S. (1992) Effects of pH and Ca^{2+} on Monomer-Dimer and Monomer-Tetramer Equilibria of Chromogranin A, *J. Biol. Chem.* 267, 11236-11241.
- Yoo, S. H. and Albanesi, J. P. (1991) High Capacity, Low Affinity Ca^{2+} Binding of Chromogranin A: Relationship Between the pH-Induced Conformational Change and Ca^{2+} Binding Property, *J. Biol. Chem.* 266, 7740-7745.




Figure 1




Figure 2

B





Figure 3




Figure 4